



Contents lists available at ScienceDirect

## International Journal of Infectious Diseases

journal homepage: [www.elsevier.com/locate/ijid](http://www.elsevier.com/locate/ijid)

# Incidental Tuberculosis in sudden, unexpected, and violent deaths in the community Lusaka, Zambia - A descriptive forensic post-mortem examination study☆☆

Luchenga Adam Mucheleng'anga<sup>1,4,11,\*</sup>, Cordilia M. Himwaze<sup>1,2,11</sup>, Viktor Telendiy<sup>1</sup>, Suzyani Simumba<sup>1</sup>, Jonathan Soko<sup>1,4</sup>, Nathan Kayonde<sup>1,2,4</sup>, Bwalya Mulenga<sup>1,2,4</sup>, Amos Hamukale<sup>1,3</sup>, Aaron Lunda Shibemba<sup>2</sup>, Patrick S. Lungu<sup>5,11</sup>, John Tembo<sup>6,11</sup>, Matthew Bates<sup>6,11</sup>, Pascalina Chanda-Kapata<sup>7</sup>, Peter Mwaba<sup>8,11</sup>, Nathan Kapata<sup>7,11</sup>, Francine Ntouni<sup>9,11</sup>, Alimuddin Zumla<sup>10,11</sup>

<sup>1</sup> Office of the State Forensic Pathologist, Zambia Ministry of Home Affairs, Lusaka Zambia

<sup>2</sup> Department of Pathology and Microbiology, University Teaching Hospital, Lusaka, Zambia

<sup>3</sup> Zambia Field Epidemiology, Lusaka, Zambia

<sup>4</sup> UNZA School of Medicine, Lusaka, Zambia

<sup>5</sup> National TB and Leprosy Programme, Ministry Of Health, Lusaka, Zambia

<sup>6</sup> HERPEZ, University Teaching Hospital, Lusaka, Zambia

<sup>7</sup> Zambia National Public Health Institute, Ministry of Health

<sup>8</sup> Apex Medical University Limited

<sup>9</sup> Fondation Congolaise pour la Recherche Médicale (FCRM), Brazzaville, Republic of Congo; Faculty of Sciences and Technology, University Marien Ngouabi, Brazzaville, Republic of Congo, University of Tübingen, Tübingen, Germany

<sup>10</sup> Division of Infection and Immunity, Center for Clinical Microbiology, University College London, and NIHR Biomedical Research Centre, UCL Hospitals NHS Foundation Trust, London, United Kingdom

<sup>11</sup> PANDORA-ID-NET Pathogenesis Group, and UNZA-UCLMS Research and Training Program\*. University Teaching Hospital, Lusaka, Zambia

## ARTICLE INFO

## Article history:

Received 7 February 2022

Revised 28 February 2022

Accepted 1 March 2022

Available online xxx

## Keywords:

Tuberculosis

Autopsy

Forensic

Incidental TB

World TB Day

## ABSTRACT

**Objectives:** Tuberculosis remains a global emergency. In Zambia only 55% of tuberculosis cases are diagnosed. We performed a study to determine incidental cases of tuberculosis seen at forensic autopsy of individuals who died suddenly and unexpectedly in the community in Lusaka, Zambia.

**Methods:** Whole-body autopsies were performed according to Standard Operating Procedures. Representative samples obtained from relevant organs were subjected to pathological examination. Information on circumstances surrounding the death was obtained. Data on patient demographics, gross and microscopic pathological findings, and cause(s) of death were analysed.

**Results:** Incidental tuberculosis was found in 52 cases (45 male, 7 female, age range 14–66) out of 4286 whole-body autopsies. 41/52 (80%) were aged 21–50 years. One was a 14-year old boy who died during a

\* Author contributions: Luchenga Adam Mucheleng'anga, Viktor Telendiy and Cordilia Maria Himwaze performed the forensic autopsy investigations. Luchenga Mucheleng'anga, Cordilia Maria Himwaze and Alimuddin Zumla ideated the study and contributed equally. Amos Hamukale analysed the data. All authors contributed to intellectual discussions and manuscript writing.

☆☆ Author declarations: All authors have a specialist interest in autopsies in Africa, TB and ONE-HEALTH. All authors declare no conflicts of interest

\* Corresponding author: Luchenga Mucheleng'anga: Ministry of Home Affairs, Office of the State Forensic Pathologist, and UNZA-UCLMS Research and Training Program, University Teaching Hospital, Lusaka, Zambia.

E-mail addresses: [luchengam@gmail.com](mailto:luchengam@gmail.com) (L.A. Mucheleng'anga), [cordeliachimwaze@gmail.com](mailto:cordeliachimwaze@gmail.com) (C.M. Himwaze), [telendijok@yandex.com](mailto:telendijok@yandex.com) (V. Telendiy), [simumbasuzyani30@gmail.com](mailto:simumbasuzyani30@gmail.com) (S. Simumba), [jonathan23soko@gmail.com](mailto:jonathan23soko@gmail.com) (J. Soko), [nathankayonde@gmail.com](mailto:nathankayonde@gmail.com) (N. Kayonde), [bmulenga1000@gmail.com](mailto:bmulenga1000@gmail.com) (B. Mulenga), [hamukalea@yahoo.com](mailto:hamukalea@yahoo.com) (A. Hamukale), [shibemba@yahoo.co](mailto:shibemba@yahoo.co) (A.L. Shibemba), [lungupatrick99@gmail.com](mailto:lungupatrick99@gmail.com) (P.S. Lungu), [john.tembo@gmail.com](mailto:john.tembo@gmail.com) (J. Tembo), [MBates@lincoln.ac.uk](mailto:MBates@lincoln.ac.uk) (M. Bates), [pascykapata@gmail.com](mailto:pascykapata@gmail.com) (P. Chanda-Kapata), [pbmwaba2000@gmail.com](mailto:pbmwaba2000@gmail.com) (P. Mwaba), [nkapata@gmail.com](mailto:nkapata@gmail.com) (N. Kapata), [fntoumi@fcrm-congo.com](mailto:fntoumi@fcrm-congo.com) (F. Ntouni), [a.zumla@ucl.ac.uk](mailto:a.zumla@ucl.ac.uk) (A. Zumla).

<https://doi.org/10.1016/j.ijid.2022.03.005>

1201-9712/© 2022 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Please cite this article as: L.A. Mucheleng'anga, C.M. Himwaze, V. Telendiy et al., Incidental Tuberculosis in sudden, unexpected, or violent deaths in the community Lusaka, Zambia - A descriptive forensic post-mortem examination study, International Journal of Infectious Diseases, <https://doi.org/10.1016/j.ijid.2022.03.005>

football match. 39/52 (75%) deaths were attributable specifically to tuberculosis only. Other deaths were due to acute alcohol intoxication(4), violence(7), ruptured ectopic pregnancy(1), bacterial meningitis (1). All the cases were from poor socio-economic backgrounds and lived in high-density areas of Lusaka.

**Conclusions:** Incidental cases of active tuberculosis undiagnosed antemortem seen at forensic autopsy reflects major gaps in the national TB control programs. More investments into proactive screening, testing, treatment activities, and accurate data collection are required.

© 2022 The Author(s). Published by Elsevier Ltd on behalf of International Society for Infectious Diseases.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

## INTRODUCTION

Community (sudden, unexpected, and violent) deaths have significant medicolegal importance (Spitz, 2006; Inquests Act). The coroner thus scrutinises these deaths with the aid of the Office of the State Forensic Pathologist (OSFP) to establish the cause and manner of death using forensic procedures (Inquests Act; The National Forensic Act 2020). Forensic procedures, specifically the forensic autopsy, provides a unique opportunity to study a range of underlying communicable and non-communicable diseases not detected ante-mortem (Mucheleng'anga et al., 2021; Himwaze et al., 2021).

Despite treatment being available, Tuberculosis (TB) remains a global emergency and is the second commonest infectious disease worldwide causing death (WHO, 2021). In 2018, the total burden of TB in Zambia was estimated to be 72 495, and of these, 40 176 (55%) were diagnosed with TB. Of the 36 431 (50%) started on treatment, only 32 700 (45%) completed treatment (Lungu et al., 2021). Forensic autopsy data may thus provide insight into the undiagnosed and sub-clinical burden of clinically undiagnosed TB in the community and who may be succumbing to the disease. Autopsies can also play a critical role in defining cryptic presentations of TB (Rampatige & Gilks, 2015). Undetected TB has significant implications as infected individuals may be a source of transmission in the community. However, the burden of undiagnosed TB in community deaths has not been estimated in Lusaka, Zambia.

We performed a case series study of incidental TB cases seen at forensic autopsy of individuals who died suddenly, unexpectedly, and violently in the community in Lusaka, Zambia. We also reviewed literature on tuberculosis findings at autopsy.

## MATERIALS AND METHODS

### Ethics approval

The coroner gave authority to conduct the forensic autopsies through an Order for Post-mortem Examination of community deaths. Forensic autopsies are mandated by law; thus, no consent or ethical permission is required to conduct the autopsy. Approval was obtained from the OSFP to access this anonymized data. The OSFP granted permission to access autopsy reports (autopsy findings and causes of death) and publish the data (Mucheleng'anga et al., 2021).

### Study design

We conducted retrospective case series analyses of whole-body forensic autopsies we had performed on community deaths in Lusaka, Zambia from January 2016 to December 2021 in which TB was an incidental finding or a cause of death. Information on the circumstances surrounding the death, symptoms, and past medical history was obtained from the next of kin by the Forensic Pathologist (FP) or Anatomic Pathologist (AP) as per procedure before the

autopsy was conducted and the Coroners Order for Post-Mortem Examination.

### Autopsy procedures

Autopsies were performed by either an FP or an AP. Personal Protective Equipment (PPE) was used, including a full gown, plastic apron, gloves (including cut-proof under gloves), face visor, boots, and N95 masks. Autopsies were performed in line with guidelines in the Practice Manual for Medicolegal Death Investigations at the OSFP in Zambia. In all cases, we followed universal precautions using PPE. A double pair of standard disposable surgical latex gloves in addition to cut-resistant gloves were also used (Mucheleng'anga, et al., 2021).

### Tissues sampled and histological examination

Representative samples were obtained from the relevant organs as required, submitted in standard tissue cassettes, and fixed in 10% neutral buffered formalin for 72 hours. The samples were processed, embedded in paraffin, sectioned, mounted onto glass slides, and stained using hematoxylin and eosin (H&E) staining according to the Standard Operating Procedure Manual at the UTH histopathology Laboratory. All slides were examined together by the FP and AP (Mucheleng'anga et al., 2021). Slides that revealed granulomatous inflammation were further subjected to Ziehl-Neelsen (ZN) staining techniques to confirm the presence of acid-fast bacilli.

### Diagnosis of Tuberculosis

The diagnosis of Pulmonary TB (PTB) was based on autopsy findings of consolidation or cavitation in the apices of the lungs, caseating lesions, histology showing granulomatous inflammation, and confirmation of tubercle bacilli by using ZN stain. Diagnosis of Extra Pulmonary TB (EPTB) was based on caseating lesions, histology showing granulomatous inflammation, and confirmation of tubercle bacilli using ZN stain.

### Data collection and analyses

Data on the decedent demographics, history, circumstances, autopsy findings, and opinion of the cause of death was entered in Excel and analysed using STATA version 14. The variables were grouped and presented as frequencies and percentages (Mucheleng'anga, et al., 2021).

### Cause(s) of Death Formulation

The cause of death was formulated within the context of the circumstances surrounding the death, history of the case, post-mortem examination findings, and ancillary studies (Mucheleng'anga et al., 2021).

**Table 1**  
Decedent's demographics, circumstances of death.

	Frequency	Percent	Cum.
<b>Age in Years</b>			
11-20	3	5.77	5.77
21-30	12	23.08	28.85
31-40	20	38.46	67.31
41-50	9	17.31	84.62
51-60	4	7.69	92.31
61-70	4	7.69	100
<b>Sex</b>			
Female	7	13.5	13.5
Male	45	86.5	100
<b>Circumstances of Death</b>			
Assaulted	4	7.69	7.69
Road Traffic Accident	3	5.77	13.46
Sudden Unexpected Death at Home	37	71.15	84.62
Sudden Unexpected Death at Work	3	5.77	90.38
Sudden Unexpected Death on Way to Hospital	5	9.62	100

**Table 2**  
Organs affected by Tuberculosis at Autopsy.

	Frequency	Percent	Cum.
Abdominal and pelvic TB	1	1.92	1.92
Cardiac and hepatic TB	1	1.92	3.85
Kaposi's sarcoma and Pulmonary TB	1	1.92	5.77
Mesenteric and Hilar lymph node TB	2	3.85	9.62
Mesenteric lymph node, Splenic and Pulmonary TB	6	11.5	19.23
Pulmonary TB	36	69.23	88.46
Pulmonary, Abdominal and Renal TB	1	1.92	90.38
Pulmonary TB, Pulmonary aspergillosis, Bacterial Meningitis, Neurocysticercosis	1	1.92	92.31
Pulmonary, Renal, Splenic and Hepatic TB	1	1.92	94.23
Pulmonary and TB Pericarditis, Cardiac cysticercosis	1	1.92	98.08
TB Pancarditis	1	1.92	100.00

## RESULTS

**Table 1** summarises the demographics and circumstances decedents. They all died suddenly, unexpectedly, and violently and were brought in dead (BID). There were fifty-two cases with active TB disease of 4286 whole-body autopsies performed. None of them had been detected or suspected ante-mortem. All decedents were from poor social-economic backgrounds and lived in high-density areas of Lusaka.

Forty-five were male, and seven were female, and the age ranged from 14-66 years. There was one paediatric TB death. The age ranges 31-40, 21-30, and 41-50 years were the most common at 20 (38.48%), 12 (23.08%), and 9 (17.31%), respectively. The age ranges 51-60 and 61-70 had 4 cases each at 7.69%, respectively. Three (5.77%) cases were in the age range 11-20.

Thirty-seven (71.15%) cases died at home, while five (9.62%) died enroute to the hospital, with three (5.77%) cases died at the workplace. Seven cases were referred for a forensic autopsy for violent deaths, three (5.77%) for road traffic accidents, and four (7.69%) for assault. One paediatric TB death involved a 14-year old boy who died suddenly while playing football.

**Table 2** summarises the autopsy findings. PTB was the most common finding in 36 (69.23%). "Mesenteric lymph node, splenic, and PTB" were present in 6 (11.5%) of cases. "Mesenteric lymph node hilar lymph node TB" was found in 2 (3.85%) cases. "Abdominal and pelvic TB," "Cardiac and hepatic TB," "Kaposi's sarcoma and PTB," "Pulmonary, abdominal, and renal TB," "PTB, aspergillosis, bacterial meningitis, neurocysticercosis," "Pulmonary, renal, splenic and hepatic TB," "Pulmonary, TB pericarditis, and Cardiac cysticercosis" and "TB pericarditis" were found at 1 case (1.92%), in each combination respectively. PTB was also found in cases with other sites of involvement in nine (9) cases (17.30%). Mesenteric lymph node involvement was found in 8 (15.35%) cases

in cases with other TB sites. Hepatic TB was found in 2 (3.85%) of cases with other TB sites. TB infection of the heart was found in 2 (3.85%) cases.

**Table 3** summarises the Cause of Death (CoD) statements. PTB and disseminated TB were the commonest causes of death in 23 (44.23%) and 10 (19.23%) cases, respectively. Acute alcohol intoxication and EPTB were CoDs in 8 cases at 7.69% each. Road traffic accidents caused three (5.76%) deaths. Blunt impact trauma due to assault, manual strangulation, and pulmonary haemorrhage due to PTB were CoDs in 6 cases at 3.85%, respectively. Two cases had CoDs attributed to ruptured ectopic pregnancy and bacterial meningitis at 1.92%, each.

### Gross pathology and microscopic findings

**Figure 1** shows representative gross pathology and microscopic pathology images.

### Gross pathology

The image in **Figure 1 A** shows an apical cavity with whitish central caseation. The image in **Figure 1 B** shows hilar lymphadenopathy with a firm consistency and grey colour **Figure 1 C** depicts Miliary pulmonary and splenic disease, with individual lesions that are small, firm (2-mm) foci of white consolidation scattered through the lung and spleen parenchyma. The images in **Figure 1 D** depicts TB pericarditis with pericardial thickening caused by fibrin and collagen formation.

### Micrographs

The micrograph in **Figure 1 E** shows a necrotic area that appears as a structureless collection of fragmented cells and amorphous granular debris within a distinctive inflammatory border in

**Table 3**  
Autopsy Cause(s) of Death.

	Frequency	Percent	Cum.
Acute Alcohol Intoxication	4	7.69	7.69
Blunt Impact trauma to head due to Assault	2	3.85	11.54
Blunt impact trauma to the head due to RTA	3	5.76	17.31
Disseminated TB	10	19.23	36.54
Ruptured Ectopic pregnancy	1	1.92	38.46
Extrapulmonary TB	4	7.69	46.15
Manual Strangulation	2	3.85	50.00
Bacterial Meningitis	1	1.92	51.92
Pulmonary TB	23	44.23	96.15
Pulmonary haemorrhage due to Pulmonary TB	2	3.85	100.00



**Figure 1.** Secondary tuberculous-apical cavitation in lung.

the lung parenchyma [Figure 1](#). F shows a necrotic area that appears as a structureless collection of lysed cells and amorphous granular debris within a distinctive border within the kidney parenchyma.

[Figures 2–6](#)

## DISCUSSION

There are several notable findings from this series. First, the majority of TB deaths occurred at home or in community settings. Second, forensic autopsies of community deaths can reveal undetected TB. Third, forty-seven of the fifty-two cases showed pulmonary disease. Fourth, PTB was a definitive cause of death in thirty-six cases and a contributing cause in eleven others. Fifth, all fifty-two cases of decedents with TB seen at autopsy were not suspected of having TB prior to death and the diagnosis was only apparent on autopsy.

Tuberculosis is the second most common cause of death from infectious diseases worldwide ([WHO, Fact sheets 2021](#)). In sub-Saharan Africa, poor economic and social factors together with suboptimal performance of national TB programs due to lack of resources required to proactively screen, find and treat all cases of TB, continue to drive the TB epidemic ([WHO, Fact sheets 2021](#)). In 2019, the incidence rate of TB was 333 per 100 000 per year in Zambia, with 15 400 TB-related deaths ([Lungu, et al., 2021](#)).



**Figure 2.** Tuberculous hilar lymphadenopathy.


Several studies conducted worldwide have shown TB at forensic autopsies. A study from India showed a 5.1% rate of TB at forensic autopsies. An ante-mortem diagnosis was not made in 84.6% ([Punia, et al., 2012](#)). Another study from Turkey reported a 1 % TB rate among forensic autopsies ([Ozsoy, et al., 2010](#)). A study in New Zealand identified TB as a cause of death in 0.2% of autopsies done ([Lum & Koelmeyer, 2005](#)). In Cape Town, South Africa, a study revealed a TB prevalence of 6.2% in persons with sudden unexpected death ([Osman et al., 2021](#)). The variation in the prevalence of TB cases identified at forensic autopsy may allude to regional differences in the prevalence of the infection.

Undiagnosed tuberculosis identified at forensic autopsy is not uncommon since TB is a chronic disease and take several weeks or months to manifest clinically ([Rastogi et al., 2011](#)). Previous autopsy studies from Zambia showed an undiagnosed case load of TB in adults and children inpatients ([Chintu, et al., 2002](#); [Bates, et al., 2015](#); [Bates et al., 2016](#); [Hussain et al., 2015](#)).

This case series shows decedents at autopsy without suspicion of having TB prior to death. Their demographics, circumstances of death, co-infections, and the extent of the disease have been highlighted. All the cases in the series were of poor social-economic status and lived in high-density areas of Lusaka. A high proportion of low socio-economic status individuals live in overcrowded households, which directly fosters TB transmission ([WHO, 2021](#)). We postulate that groups from high-density and poor social-economic status areas, household contacts of those with infectious TB, and contacts known to be HIV positive must be targeted for screening given that Lusaka high-density areas are also known to have high HIV prevalence.

Forty-one of fifty-two cases were in the age range 21–50 years, with the 31–40 range contributing most cases. This find-







**Figure 3.** Miliary tuberculosis in Lung (Left) and Spleen (Right).



**Figure 4.** Fibrinous tuberculous pericarditis.



**Figure 5.** Caseating granuloma in lung (Micrograph at x 4 magnification).



**Figure 6.** Caseating granuloma in kidney (Micrograph at magnification x 10).

ing contrasts with a study in Germany which showed that most of the cases were within the age ranges of 30–59 and 80–99 (Theegarten, et al., 2006). The difference is related to the differences in HIV prevalence and the social-economic status of the countries. The difference may also be attributed to differences in the age structure of the two populations (Mucheleng'anga, et al., 2021). However, our results are similar to a study in a forensic population by Himwaze et al., 2020 in Lusaka, who found similar results and concluded that this age group was most at risk of premature death due to increased alcohol intake and outdoor activities in

the forensic population (Himwaze, et al., 2020). There were more males than females in our series. This is consistent with our previous autopsy studies (Mucheleng'anga, et al., 2021; Himwaze, et al., 2020). It has been suggested that males are more predisposed to sudden, unexpected, and violent deaths due to complications of disease arising from poor health-seeking habits (Himwaze, et al., 2020).

Regarding the circumstances surrounding the death, many sudden and unexpected deaths occurred outside the hospital facility, either at home, at work, or on the way to the health facility in the

series. These circumstances again emphasize the hypothesis that this predominantly male population dies suddenly and unexpectedly outside the health facility due to complications of disease arising from poor health-seeking habits (Mucheleng'anga, et al., 2021; Himwaze, et al., 2020). An autopsy study in South Africa revealed a high proportion of pulmonary TB in people who died at home without an apparent cause of death. This was attributed to the high burden of HIV infection (Omar, et al., 2015) Himwaze, et al., 2020., found an HIV prevalence of 31% in the forensic population at autopsy in Lusaka (Himwaze, et al., 2020). The forensic population is known to suffer from advanced disease due to the self-rationalisation of symptoms and a failure to seek medical attention (Spitz, 2006; Mucheleng'anga et al., 2021). The one paediatric TB death involved a 14-year-old boy who died suddenly during a football match. This is a telling example where TB kills "silently." At autopsy, a TB pancarditis was diagnosed. The screening of TB contacts for individuals who are both symptomatic and asymptomatic, often neglected, can be an essential benefit to the community of individual children. This child would have benefited from TB treatment if screening had been conducted (Omar, et al., 2015).

Pulmonary TB (apical cavitation, consolidation, granulomatous inflammation), was our series' most common autopsy finding. These were confirmed on histology and confirmed by positive ZN staining. We observe that the spleen and mesenteric lymph nodes were the second and third most common sites of infection in this series. Three cases had coinfections, the first with aspergillosis, bacterial meningitis and neurocysticercosis. The second with Kaposi's sarcoma and HIV, while the third with TB pericarditis and cardiac cysticercosis.

TB was the primary cause of death in thirty-nine of fifty-two incidental TB cases of TB seen at autopsy, with Pulmonary (23 cases) and disseminated (10 cases) being the commonest CoD, respectively. Four cases had pulmonary TB but died of acute alcohol intoxication. Seven cases were violent deaths but revealed TB on autopsy. Evidence of undiagnosed TB suggests that TB-related mortality is under-ascertained and under reported. There is therefore need for more research into active screening methods in the community as autopsies provide the most accurate data about cause of death despite being operationally difficult and expensive. We postulate that more TB cases could be found by searching more actively among specific population groups (Omar, et al., 2015).

Diagnosis of incidental TB in community deaths reflects the burden of disease within the community and while public health efforts worldwide are being directed towards reducing the spread of TB, the forensic autopsy is a vital tool in detecting unknown TB cases in the community (Mucheleng'anga, et al., 2021). Forensic autopsy data are thus indispensable for disease intelligence and prevention and may be helpful and lead to timely medical intervention for the decedent's relatives who might have been exposed to TB. These data can help public health map disease surveillance systems, implement public awareness programs for prevention, and treat sources in households and the community (Rasika & Gilks, 2015; Dye, et al., 2006 ).

There have been previous calls to conduct forensic autopsy studies in community deaths to identify undiagnosed TB cases (Syed, et al., 2015). The actual data on the causes of deaths in the community (rural and urban) and in hospitals in most SSA countries remain unknown, and the quality of cause-specific mortality statistics remains poor. Thus, there is an urgent need to obtain more accurate cause-of-death data from SSA countries by reviving routine autopsies which declined in most SSA countries because of several reasons, such as lack of resources to maintain an effective pathology service, scarcity of trained pathologists, inadequate infrastructure, and difficulties in obtaining consent for autopsies due

to cultural beliefs and religious sensitivities (Chintu, et al., 2002). This case series is one such activity related to that clarion call.

Our data should be viewed under the limitations of the study. This case series is limited by the small number of cases because not all violent deaths such as suicides and road traffic accidents undergo a whole body autopsy, possibly leading to missed incidental TB. Additionally culture and polymerase chain reaction testing was not done. ZN staining that was used to confirm the presence of tubercle bacilli in tissue at autopsy does not distinguish tuberculous from non-tuberculous mycobacterial infections. Future autopsy studies should include mycobacterial culture and molecular analyses for more accurate diagnosis of *Mycobacterium tuberculosis* (*Mtb*) in autopsy tissues.

Our autopsy study serves as a pivotal starting point for conversations around future autopsy studies in the SSA focused on TB pathology, pathogenesis and co-infections. Undiagnosed TB cases and other co-morbidities such as HIV, malaria, COVID-19, hypertension, diabetes, and other diseases require further investigation. Our previous COVID-19 autopsy studies in Zambia have highlighted the need for sequencing data to align to autopsy studies (Mucheleng'anga, et al., 2021; Himwaze et al., 2021). Apart from more investments into TB programs, for more proactive screening, testing and treatment activities, accurate data collection, revamping autopsy studies worldwide would provide more insights into the actual mortality burden. *Mtb* can persist intracellularly in lung tissue without histological evidence of tuberculous lesions (Hernandez-Pando et al, 2000), and autopsy studies provide opportunities to study latent TB infection, early TB disease not manifest clinically and host-*Mtb* interactions.

## Declaration of Competing Interest

The authors declare no conflict of interest.

## Author contributions

Luchenga Mucheleng'anga and Alimuddin Zumla ideated the study and contributed equally. Luchenga Adam Mucheleng'anga, Viktor Telendiy and Cordilia Maria Himwaze performed the forensic autopsy investigations. Amos Hamukale analysed the data. All authors contributed to intellectual discussions and manuscript writing.

## Acknowledgements

We thank the mortuary staff for their assistance with performing the autopsies.

Authors are co-Investigators of the Pan-African Network on Emerging and Re-Emerging Infections (PANDORA-ID-NET – <https://www.pandora-id.net/>) funded by the European and Developing Countries Clinical Trials Partnership the EU Horizon 2020 Framework Programme. AZ and FN acknowledge support from EDCTP CANTAM-3. Sir Zumla is a Mahathir Science Award and EU-EDCTP Pascoal Mocumbi Prize Laureate, and is in receipt of a UK National Institutes of health rEsearch Senior Investigator Award.

## Funding

The authors did not receive any funding for this series.

## References

- Bates M, Shibemba A, Mudenda V, Chimoga C, Tembo J, Kabwe M, et al. Burden of respiratory tract infections at post mortem in Zambian children. *BMC Med* 2016;14:99. doi:[10.1186/s12916-016-0645-z](https://doi.org/10.1186/s12916-016-0645-z).



- Bates M, Mudenda V, Shibemba A, Kaluwaji J, Tembo J, Kabwe M, et al. Burden of tuberculosis at post mortem in inpatients at a tertiary referral centre in sub-Saharan Africa: a prospective descriptive autopsy study. *Lancet Infect Dis* 2015;15(5):544–51.
- Chintu C, Mudenda V, Lucas S, Nunn A, Lishimpi K, Maswahu D, et al. Lung diseases at necropsy in African children dying from respiratory illnesses: a descriptive necropsy study. *Lancet* 2002;360(9338):985–90.
- Dye C, Harries AD, Maher D, et al. Tuberculosis. In: Jamison DT, Feachem RG, Makgoba MW, et al, editors. *Disease and Mortality in Sub-Saharan Africa*. Washington (DC): The International Bank for Reconstruction and Development /The World Bank; 2006 Chapter 13. Available from <https://www.ncbi.nlm.nih.gov/books/NBK2285/>.
- Himwaze CM, Telendiy V, Maate F, Mupeta S, Chitalu C, Chanda D, et al. Post-mortem examination of Hospital Inpatient COVID-19 Deaths in Lusaka, Zambia - A Descriptive Whole-body Autopsy Series. *Int J Infect Dis* 2021;108:363–9 Epub 2021 Jun 17. PMID: 34146690; PMCID: PMC8215884. doi:[10.1016/j.ijid.2021.06.013](https://doi.org/10.1016/j.ijid.2021.06.013).
- Himazwe C, Mucheleng'anga L, Siyumbwa SN, Kaile T, Julius P. Prevalence of Human Immunodeficiency Virus, Hepatitis B, and Hepatitis C Viral Infections among Forensic Autopsy Cases at the University Teaching Hospital in Lusaka, Zambia. *Forensic Science International: Reports* 2020;2 Article 100133 <https://doi.org/10.1016/j.fsir.2020.100133>. doi:[10.1016/j.fsir.2020.100133](https://doi.org/10.1016/j.fsir.2020.100133).
- Hussain SA, Madadin M, Menezes RG. Tuberculosis burden at post mortem. *Lancet Infect Dis* 2015;15(10):1133–4 Epub 2015 Sep 20. PMID: 26461944. doi:[10.1016/S1473-3099\(15\)00307-2](https://doi.org/10.1016/S1473-3099(15)00307-2).
- Inquests Act, Vol:4 Chapter 36 of the Laws of Zambia <https://www.parliament.gov.zm/node/736>. Accessed on February 1st, 2022.
- Lum D, Koelmeyer T. Tuberculosis in Auckland autopsies, revisited. *N Z Med J* 2005;118(1211):U1356 PMID: 15778757.
- Lungu P, Kerkhoff AD, Kasapo CC, Mzyece J, Nyimbili S, Chimzizi R, et al. Tuberculosis care cascade in Zambia - identifying the gaps in order to improve outcomes: a population-based analysis. *BMJ Open* 2021;11(8). doi:[10.1136/bmjopen-2020-044867](https://doi.org/10.1136/bmjopen-2020-044867).
- Mucheleng'anga LA, Telendiy V, Hamukale A, Shibemba AL, Zumla A, Himwaze CM. COVID-19 and Sudden Unexpected Community Deaths in Lusaka, Zambia, Africa - A Medico-Legal Whole-Body Autopsy Case Series. *Int J Infect Dis* 2021;109:160–7 Epub 2021 Jul 5. PMID: 34320461; PMCID: PMC8255190. doi:[10.1016/j.ijid.2021.07.001](https://doi.org/10.1016/j.ijid.2021.07.001).
- Osman M, Verster J, Dempers JJ, Du Preez K, von Delft A, Dunbar R, et al. Tuberculosis in persons with sudden unexpected death, in Cape Town, South Africa. *Int J Infect Dis* 2021;105:75–82 Epub 2021 Feb 12. PMID: 33582368; PMCID: PMC8358423. doi:[10.1016/j.ijid.2021.02.036](https://doi.org/10.1016/j.ijid.2021.02.036).
- Omar T, Variava E, Moroe E, Billioux A, Chaisson RE, Lebina L, et al. Undiagnosed TB in adults dying at home from natural causes in a high TB burden setting: a post-mortem study. *Int J Tuberc Lung Dis* 2015;19(11):1320–5 PMID: 26467584. doi:[10.5588/ijtld.15.0222](https://doi.org/10.5588/ijtld.15.0222).
- Ozsoy S, Demirel B, Albay A, Kisa O, Dinc AH, Safali M. Tuberculosis prevalence in forensic autopsies. *Am J Forensic Med Pathol* 2010;31(1):55–7 PMID: 19935395. doi:[10.1097/PAF.0b013e3181c215f9](https://doi.org/10.1097/PAF.0b013e3181c215f9).
- Punia RS, Mundi I, Mohan H, Chavli KH, Harish D. Tuberculosis prevalence at autopsy: a study from North India. *Trop Doct* 2012;42(1):46–7 PMID: 22290110. doi:[10.1258/td.2011.110314](https://doi.org/10.1258/td.2011.110314).
- Rampatige R, Gilks CF. Autopsies and better data on causes of death in Africa. *Lancet Infect Dis* 2015;15(5):492–4 Epub 2015 Mar 10. PMID: 25765218. doi:[10.1016/S1473-3099\(15\)70096-4](https://doi.org/10.1016/S1473-3099(15)70096-4).
- Rastogi P, Kanchan T, Menezes RG. Sudden unexpected deaths due to tuberculosis: an autopsy based study. *J Forensic Med Toxicol* 2011;28:81–5.
- Spitz WU. *Medicolegal Investigation of Death: Guidelines for the Application of Pathology to Crime Investigation*. 4th edition. Charles C, Springfield; 2006 II.
- Theegarten D, Kahl B, Ebsen M. Häufigkeit und Morphologie der Tuberkulose bei Obduktionen: Zunahme aktiver Formen [Frequency and morphology of tuberculosis in autopsies: increase of active forms]. *Dtsch Med Wochenschr* 2006;131(24):1371–6 German PMID: 16783670. doi:[10.1055/s-2006-946581](https://doi.org/10.1055/s-2006-946581).
- The National Forensic Act, 2020, Act number:2. <https://www.parliament.gov.zm/node/8571>, Accessed on Feb 20th 2022.
- WHO, Fact sheets 2021, Tuberculosis. <https://www.who.int/news-room/fact-sheets/detail/tuberculosis>, accessed 3rd February, 2022
- WHO. Global tuberculosis report 2021. October 14, 2021 <https://www.who.int/publications/i/item/9789240037021>, accessed 30th January, 2022.